

Excited states of a static dilute spherical Bose condensate in a trap*

Alexander L. Fetter

Departments of Physics and Applied Physics, Stanford University, Stanford, CA 94305-4060

(February 1, 2008)

The Bogoliubov approximation is used to study the excited states of a dilute gas of N atomic bosons trapped in an isotropic harmonic potential characterized by a frequency ω_0 and an oscillator length $d_0 = \sqrt{\hbar/m\omega_0}$. The self-consistent static Bose condensate has macroscopic occupation number $N_0 \gg 1$, with nonuniform spherical condensate density $n_0(r)$; by assumption, the depletion of the condensate is small ($N' \equiv N - N_0 \ll N_0$). The linearized density fluctuation operator $\hat{\rho}'$ and velocity potential operator $\hat{\Phi}'$ satisfy coupled equations that embody particle conservation and Bernoulli's theorem. For each angular momentum l , introduction of quasiparticle operators yields coupled eigenvalue equations for the excited states; they can be expressed either in terms of Bogoliubov coherence amplitudes $u_l(r)$ and $v_l(r)$ that determine the appropriate linear combinations of particle operators, or in terms of hydrodynamic amplitudes $\rho'_l(r)$ and $\Phi'_l(r)$. The hydrodynamic picture suggests a simple variational approximation for $l > 0$ that provides an upper bound for the lowest eigenvalue ω_l and an estimate for the corresponding zero-temperature occupation number N'_l ; both expressions closely resemble those for a uniform bulk Bose condensate.

PACS numbers: 03.75.Fi, 05.30.Jp, 32.80.Pj, 67.90.+z

*Contributed paper for Low Temperature Conference LT21, Prague, August, 1996, to appear in Czechoslovak Journal of Physics

I. INTRODUCTION

Recent experimental verification of Bose condensation in dilute confined ^{87}Rb [1] has stimulated extensive theoretical research. In the Bogoliubov approximation, N_0 atoms occupy the macroscopic condensate, with $N_0 \lesssim N$ and only small depletion ($N'/N \equiv 1 - N_0/N \ll 1$). For a repulsive scattering length $a > 0$ and mean atomic density n , this depletion is of order $\sqrt{na^3}$ [2] and thus nonperturbative.

The balance between kinetic energy $\hbar^2/2m\xi^2$ and interaction energy $4\pi a\hbar^2 n/m$ defines the “healing length” $\xi = (8\pi an)^{-1/2}$, and the Bogoliubov approximation requires that $a \ll \xi$. In addition, a confined Bose gas with typical dimension R differs qualitatively from an ideal Bose gas whenever $\xi \ll R$. The present work uses the Bogoliubov approximation to provide a hydrodynamic description of the excited states and a variational estimate of the eigenvalue ω_l and zero-temperature occupation number N'_l for the lowest radial mode for each positive angular momentum $l \geq 1$.

II. BASIC FORMALISM

In [1], the harmonic trap $U = \frac{1}{2}m\omega_0^2 r^2$ (here taken as isotropic) has a size $d_0 = \sqrt{\hbar/m\omega_0} \approx 1 \mu\text{m}$, and the positive scattering length $a \approx 10 \text{ nm}$ acts to expand the atomic cloud. For $\mathcal{N} \equiv Na/d_0 \gg 1$, the actual atomic density $\approx N/R^3$ is smaller than the naive estimate $\approx N/d_0^3$, with the radial expansion factor R/d_0 of order $\mathcal{N}^{1/5}$ [3]; thus the Bogoliubov approximation holds for $Na^3/R^3 \approx \mathcal{N}^{2/5}(a/d_0)^2 \ll 1$ [namely, $N \ll (d_0/a)^6 \sim 10^{12}$]. The additional condition $R \gg \xi$ for the failure of an ideal-Bose-gas model now requires $(8\pi\mathcal{N}^{4/5})^{1/2} \gg 1$, which here holds even for $N \approx 100$.

The condensate wave function Ψ satisfies the nonlinear Gross-Pitaevskii equation [4,5]

$$(T + U - \mu + V) \Psi = 0, \quad (1)$$

where $T = -\hbar^2 \nabla^2 / 2m$ is the kinetic energy, $V \equiv 4\pi a\hbar^2 m^{-1} |\Psi|^2$ characterizes the repulsive interaction energy, and the chemical potential μ must be adjusted to ensure that $\int dV |\Psi|^2 =$

N_0 and that $\Psi \rightarrow 0$ at large distances. In the present case, the condensate is stationary with density $n_0(r) = |\Psi(r)|^2$. For large $\mathcal{N}$, the kinetic energy can be ignored [3], giving the “Thomas-Fermi” approximation

$$V_{TF} = (\mu - U) \theta(\mu - U) \equiv \frac{1}{2} \hbar \omega_0 (R^2 - r^2) \theta(R - r), \quad (2)$$

where lengths are expressed in units of d_0 , $\mu = \frac{1}{2} \hbar \omega_0 R^2$ defines the dimensionless “radius” R of the condensate, and $R^5 = 15\mathcal{N}$ [3].

III. HYDRODYNAMIC DESCRIPTION

The (small) noncondensate field operator $\hat{\phi}$ obeys the linear equation

$$i\hbar \partial \hat{\phi} / \partial t = (T + U - \mu + 2V) \hat{\phi} + V \hat{\phi}^\dagger, \quad (3)$$

along with the adjoint equation. In the Bogoliubov approximation ($N' \ll N_0$), the total particle density operator becomes $\hat{\psi}^\dagger \hat{\psi} \approx n_0 + \hat{\rho}'$, where the operator $\hat{\rho}' = \sqrt{n_0}(\hat{\phi} + \hat{\phi}^\dagger)$ characterizes the fluctuating noncondensate density; similarly, the particle current operator here becomes $\hat{\mathbf{j}}' \approx n_0 \nabla \hat{\Phi}'$, where $\hat{\Phi}' = (\hbar/2mi\sqrt{n_0})(\hat{\phi} - \hat{\phi}^\dagger)$ is the velocity potential operator. Equation (3) readily yields

$$\partial \hat{\rho}' / \partial t + \nabla \cdot (n_0 \nabla \hat{\Phi}') = 0, \quad (4)$$

and the linearized Bernoulli’s equation [6]

$$\frac{\partial \hat{\Phi}'}{\partial t} + \frac{4\pi a \hbar^2}{m^2} \hat{\rho}' + \frac{\hbar^2}{4m^2 n_0} \left[\nabla \cdot \left(\hat{\rho}' \frac{\nabla n_0}{n_0} \right) - \nabla^2 \hat{\rho}' \right] = 0. \quad (5)$$

The latter can be recognized as a linearization of the exact classical Bernoulli’s theorem [7] $U + \frac{1}{2}mv^2 + (e + p)/n + m\partial\Phi/\partial t = 0$ for a compressible irrotational isentropic fluid with density $n_0 + \hat{\rho}'$ and velocity potential $\hat{\Phi}'$ [here, $p(n) = 2\pi a \hbar^2 n^2/m$ is the pressure functional and $e(n) = \sqrt{n} T \sqrt{n} + p(n)$ is the (quantum) energy-density functional].

IV. BOGOLIUBOV EQUATIONS

The hydrodynamic description provides a valuable qualitative picture of the normal modes of the nonuniform compressible spherical condensate, but the resulting equations involve spatial derivatives of n_0 ; furthermore, in the Thomas-Fermi approximation (2), the appropriate boundary conditions at R are not obvious. Thus it is preferable to return to Eq. (3), expressing the field operator $\hat{\phi}$ as a linear combination of quasiparticle operators α_j and $\alpha_j^\dagger$

$$\hat{\phi}(\mathbf{r}, t) = \sum_j' \left[u_j(\mathbf{r}) \alpha_j e^{-i\omega_j t} - v_j^*(\mathbf{r}) \alpha_j^\dagger e^{i\omega_j t} \right]; \quad (6)$$

here, the wave-function coefficients u_j and v_j constitute a two-component vector function $\mathcal{U}_j$ that obeys the matrix “Bogoliubov” eigenvalue equation [6]

$$(T + U - \mu + 2V) \mathcal{U}_j - V \tau_1 \mathcal{U}_j = \hbar \omega_j \tau_3 \mathcal{U}_j. \quad (7)$$

In contrast to Eqs. (4) and (5), only $V \propto n_0$ appears here, and the quantum interpretation provides the obvious boundary condition that $\mathcal{U}_j$ vanish for $r \rightarrow \infty$ (note that the hydrodynamic amplitudes are essentially linear combinations of u_j and v_j).

In the case of a static spherical condensate, the Bogoliubov Eqs. (7) have solutions of the form $u_l(r)Y_{lm}$ and $v_l(r)Y_{lm}$; in addition, they have a simple variational basis [6] that yields an upper bound for the lowest eigenvalue ω_l for each positive angular momentum $l > 0$. To ensure the proper normalization, take $u_l(r) = \cosh \chi_l f_l(r)$ and $v_l(r) = \sinh \chi_l f_l(r)$, where $\int_0^\infty r^2 dr |f_l(r)|^2 = 1$. With the Thomas-Fermi approximation (2), the analogy to acoustic waves in a sphere suggests the radial trial function $f_l(r) \propto x^l(1 - x^2)\theta(1 - x)$, where $x \equiv r/R$. A simple variational calculation yields the estimate $\omega_l/\omega_0 = \sqrt{T_l^2 + 2T_l V_l}$, with $T_l = \frac{1}{4}(2l + 3)(2l + 7)/R^2$ and $V_l = 3R^2/(2l + 9)$ the expectation values of the dimensionless kinetic energy $-\frac{1}{2}\nabla^2$ and Thomas-Fermi potential energy $\frac{1}{2}(R^2 - r^2)$. This frequency has a very different form for small and large l

$$\frac{\omega_l}{\omega_0} \approx \begin{cases} \sqrt{\frac{3}{2}(2l + 3)(2l + 7)/(2l + 9)}, & \text{for } T_l \ll V_l; \\ (2l + 3)(2l + 7)/4R^2, & \text{for } T_l \gg V_l. \end{cases} \quad (8)$$

Stringari [8] has obtained a related expression $\omega_l/\omega_0 = \sqrt{l}$ with a purely hydrodynamic model, adding that $\omega_1 = \omega_0$ is an exact result. Note the close analogy to the Bogoliubov excitation energy $E_k = \sqrt{T_k^2 + 2T_k V_k}$ for a uniform bulk medium, where $T_k = \hbar^2 k^2/2m$ and $V_k = 4\pi a \hbar^2 n_0/m$ [2].

Each of these lowest excited states for $l > 0$ has a zero-temperature occupation number $N'_l = \int_0^R r^2 dr |v_l(r)|^2$, with the variational estimate $N'_l = \sinh^2 \chi_l = \frac{1}{2}(T_l + V_l)/\sqrt{T_l^2 + 2T_l V_l} - \frac{1}{2}$, again very similar in structure to that for a uniform Bose gas [2]. In particular, $N'_l \approx \sqrt{V_l/8T_l} \gg 1$ for $T_l \ll V_l$, and $N'_l \approx V_l^2/4T_l^2 \ll 1$ for $T_l \gg V_l$.

ACKNOWLEDGMENTS

Supported in part by the National Science Foundation under Grant No. DMR 94-21888; I am grateful for helpful correspondence with B. V. Svistunov.

-
- [1] M. H. Anderson *et al.*, Science **269**, 198 (1995).
 - [2] See, for example, A. L. Fetter and J. D. Walecka, *Quantum Theory of Many-Particle Systems* (McGraw-Hill, NY, 1971), Secs. 35 and 55.
 - [3] G. Baym and C. J. Pethick, Phys. Rev. Lett. **76**, 6 (1996).
 - [4] E. P. Gross, Nuovo Cimento **20**, 451 (1961).
 - [5] L. P. Pitaevskii, Zh. Eksp. Teor. Fiz. **40**, 646 (1961); Sov. Phys.-JETP **13**, 451 (1961).
 - [6] A. L. Fetter, Phys. Rev. A, to be published.
 - [7] See, for example, A. L. Fetter and J. D. Walecka, *Theoretical Mechanics of Particles and Continua* (McGraw-Hill, NY, 1980), Sec. 48.
 - [8] S. Stringari, unpublished (cond-mat/9603126).